

AD\_\_\_\_\_

Award Number: DAMD17-00-1-0384

TITLE: Solid-Phase Combinatorial Approach to Estradiol  
Tamoxifen/Raloxifene Hybrids: Novel Chemotherapeutic/  
Prophylactic Selective Estrogen Receptor Modulators  
(SERM)

PRINCIPAL INVESTIGATOR: Robert N. Hanson, Ph.D.

CONTRACTING ORGANIZATION: Northeastern University  
Boston, Massachusetts 02115-5000

REPORT DATE: July 2001

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20020311 140

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> July 2001	<b>3. REPORT TYPE AND DATES COVERED</b> Annual (1 Jul 00 - 30 Jun 01)	
<b>4. TITLE AND SUBTITLE</b> Solid-Phase Combinatorial Approach to Estradiol Tamoxifen/Raloxifene Hybrids: Novel Chemotherapeutic/ Prophylactic Selective Estrogen Receptor Modulators (SERM)			<b>5. FUNDING NUMBERS</b> DAMD17-00-1-0384	
<b>6. AUTHOR(S)</b> Robert N. Hanson, Ph.D.				
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Northeastern University Boston, Massachusetts 02115-5000  E-Mail: r.hanson@neu.edu			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			<b>10. SPONSORING / MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b>				
<b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for Public Release; Distribution Unlimited.				<b>12b. DISTRIBUTION CODE</b>
<b>13. ABSTRACT (Maximum 200 Words)</b>  The objective of this project is the development of new chemotherapeutic agents for the treatment of hormone-responsive breast cancer using a solid phase approach to synthesize new agents having features common to both steroids and antiestrogens. During the past year we have functionalized the carboxy resin with both the E- and Z-tributylstannylvinyl estradiol. We have prepared and characterized the initial series of iodophenoxyalkylamines that will be coupled to the resin-bound steroid. The biological assays have been established and validated using series of previously prepared compounds. Molecular modeling and NMR spectrometric methods for conformational analyses and for data analysis have been developed. The next years specific aims involve the synthesis and evaluation of the initial series of target compounds.				
<b>14. SUBJECT TERMS</b> Breast Cancer				<b>15. NUMBER OF PAGES</b> 46
				<b>16. PRICE CODE</b>
<b>17. SECURITY CLASSIFICATION OF REPORT</b> Unclassified	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> Unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> Unclassified	<b>20. LIMITATION OF ABSTRACT</b> Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)  
Prescribed by ANSI Std. Z39-18  
298-102

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

X Where copyrighted material is quoted, permission has been obtained to use such material.

    Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

    Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

N/A For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

  
P.I.

7 Dec. 2001

## **Table of Contents**

1. Cover	1
2. SF 298	2
3. Foreword	3
4. Table of Contents	4
5. Introduction	5
6. Body	6-9
7. Key Research Accomplishments	10
8. Reportable Outcomes	11
9. Conclusions	11
10. References	12
11. Appendix	12

## **5. Introduction**

The overall objective of this project is the development of new chemotherapeutic agents for the treatment or prevention of hormone-responsive breast cancer. Based upon our previous synthetic work and molecular modeling studies, we undertook the preparation of a series of 17 $\alpha$ -(substituted phenylvinyl) estradiols using a solid-phase synthesis approach. The key feature of these compounds is the incorporation onto the phenylvinyl group of the dialkylaminoalkoxy side chain found in the potent anti-estrogens tamoxifen and raloxifene. This would be accompanied by the development of the appropriate biological assays to determine bioactivity and NMR evaluations to determine the molecular basis for the observed effects. In this report we describe the preliminary aspects of this work and the advances that we have made in these areas.

## **6. Body**

The research proposal described 5 Tasks as part of the Statement of Work. These included: 1. Initial target compound design. 2. Chemical synthesis of target compounds in initial directed library. 3. Measurement of biological properties-affinity and efficacy. 4. Assessment of structure-activity relationships. 5. Chemical synthesis of target compounds in second generation libraries. Work has been undertaken on the first three tasks and will be described in the report.

### **Task 1. Initial target compound design (Pre-award).**

Prior to the initiation of the award, we evaluated the SAR for the tamoxifen and raloxifene anti-estrogens and selected the structural variations for the dialkylaminoalkoxy side chains. Based upon our analysis of the crystal structures for liganded ER-LBDs, we chose to prepare the dimethylamino-, diethylamino-, pyrrolidino- and piperidino- tertiary amines with ethoxy, propoxy- and butoxy- side chains situated at the ortho-, meta- or para- positions of the phenyl ring. The starting materials were purchased from commercial sources. The methods for the biological assays were identified as well.

### **Task 2, Chemical synthesis of target compounds in initial directed library (Months 1-18).**

During this period we have focused on two aspects. The first is the preparation of the functionalized resin that will be used in the combinatorial chemistry. The second is

the preparation of the iodophenoxy precursors for the penultimate coupling step. For the first part, we have prepared on large scale the E- and Z-17 $\alpha$ -tributylstannyl vinyl estradiol isomers. While the E-isomer predominated (7.5g vs. 2.0 g), there was sufficient quantities of both to perform the coupling to the carboxy resin (polystyrene backbone). We used carbonyl diimidazole as the coupling reagent. For the second part, we used the commercially available ortho-, meta- and para-iodophenols as starting materials. These could be coupled directly with the commercially available N-hydroxyethyl amines using the Mitsunobu reaction. Preparation of the corresponding N-hydroxypropyl- and N-hydroxybutyl- derivatives was accomplished in two steps. We employed the Mitsunobu reaction to couple the phenol to the w-halopropan(butan)ol followed by amination with the desired secondary amine. The final compounds were characterized by NMR and MS.

The project is now ready to undertake the solid-phase coupling of the two components using the Stille reaction conditions that we have used in other studies.

### **Task 3. Measurement of biological properties-affinity and efficacy (Months 1-36).**

We have established the biological assay systems for evaluating the compounds and have begun to determine their properties. Our initial results examined the effect of stereochemistry (E-vs Z-) and position (o-/m-/p-) on receptor binding for the hydroxyphenyl vinyl estradiols and for the para-methoxyphenyl vinyl estradiols. In this series, the hydroxy and methoxy substituents are well tolerated. Although the binding

(RBA values) are less than estradiol, except for the E-m-hydroxyphenyl compound which is comparable to estradiol itself, they are equal to or better than the unsubstituted phenyl vinyl estradiol. Only the value for the Z-m-isomer is unavailable at this time. The uterotrophic growth assay will be done in the near future to determine the efficacy of the compounds and establish a baseline against which the dialkylaminoalkoxyphenyl products can be compared.

One key aspect in this portion of the project that was determined in one of our other programs was the need to do the in vivo assays with a single lot of animals. The lot to lot variation was too great to permit a reliable comparison of results for compounds in a series performed at different times. Therefore, we have to wait until we have all six isomers of a series (E-/Z-; o-/m-/p-) before conducting an assay.

#### **Task 4. Assessment of structure-biological activity relationships (Months 6-36).**

We have utilized both NMR (H-1;C-13) and molecular modeling to examine the conformation of the substituted phenyl vinyl estradiols. In general, the molecular modeling provides several reasonable low energy conformers for the compounds. Two of these conformers are observed in solution using NMR. However, the energy barrier between the two (and sometimes three) low energy conformers is low and it appears that small energy contributions from the receptor protein could easily induce a change to give a conformation not favored in solution or necessarily predicted by computational methods. We have undertaken the evaluation of the ER-LBD based on the several



published structures. This evaluation includes molecular dynamics with and without bound ligand. Since several of the initial compounds that we have evaluated in other studies are estrogen receptor agonists, our first studies utilized the agonist form of the receptor. As we get more information about our new ligands, we can investigate their docking modes with the antagonist form of the receptor. To date, our studies suggest that there is significant interaction between the substituted phenyl vinyl group and the amino acid sidechains present in the helix-11/12 region but definite sites cannot be identified.

## **7. Key Research Accomplishments.**

- Prepared functionalized resin with both E- and Z-triethylstannylvinyl estradiol
- Prepared many of the dialkylaminoalkoxyphenyl iodide coupling reagents
- Developed appropriate NMR and molecular modeling methods using structurally similar estradiol derivatives
- Developed biological assays and validated methods using structurally similar estradiol derivatives
- Initiated SAR studies using data from parallel series of compounds

## **8. Reportable Outcomes.**

a. Manuscripts, abstracts, presentations.

1. Sebag, A.B., Hanson, R.N., Forsyth, D.A., and Lee, C.-Y.,  
Conformational studies of novel estrogen receptor ligands by 1D and 2D  
NMR spectroscopy and computational methods. Org. Magn. Res.  
(accepted pending minor revisions).
2. Hanson, R.N., Design, synthesis and evaluation of novel  
steroidal antiestrogens for the treatment of hormone responsive breast  
cancer. Invited presentation at The Philip S. Portoghese Symposium in  
Medicinal Chemistry, August 23-24, 2001, Minneapolis, Minnesota.

a. Degrees obtained supported by the award.

None.

## **9. Conclusions.**

At this point we have achieved several of the intermediate objectives of the project. We have prepared the E- and Z-tributylstannylvinyl estradiols and coupled them to the carboxy resin. We have prepared a number of the dialkylaminoalkoxyphenyl iodides that will we used to react with the functionalized resin. Using our Stille coupling procedure we have prepared sample substituted phenylvinyl estradiol derivatives on the resin and submitted them to validate our biological assays. We have established the assay for receptor affinity using the ER-LBD overexpressed in the BL-21 cell line. We are

working on the MCF-7 cell proliferation assay to generate the reproducibility necessary for this project. We have established the immature rat uterotrophic growth assay and tested it using a set of estradiol derivatives made previously. These results suggest significant differences in potency exist between E- and Z-, ortho-, meta- and para-isomers. Progress made to date in all areas of the project indicate that goals for the next year are achievable.

#### **10. References:**

None.

#### **11. Appendix:**

The appendix material is comprised of copies of one manuscript accepted for publication in Organic Magnetic Resonance, pending final revisions, and the brochure for The Philip S. Portoghese Symposium in Medicinal Chemistry.

## Conformational Studies of Novel Estrogen Receptor Ligands by 1D and 2D NMR Spectroscopy and Computational Methods

Albert B. Sebag, Robert N. Hanson\*, David A. Forsyth, Choon Young Lee

*Departments of Pharmaceutical Science and Chemistry, Northeastern University, 360 Huntington Avenue, Boston, MA 02115*

**Abstract:** The solution conformations of the novel estrogen receptor ligands, (17 $\alpha$ , 20E)-(p- $\alpha$ ,  $\alpha$ ,  $\alpha$ -trifluoromethylphenyl)vinyl estradiol (**1**) and (17 $\alpha$ , 20E)-(o- $\alpha$ ,  $\alpha$ ,  $\alpha$ -trifluoromethylphenyl)vinyl estradiol (**2**) were investigated in 2D and 1D NOESY studies and by comparison of  $^{13}\text{C}$  NMR chemical shifts with theoretical shieldings. The  $^1\text{H}$  and  $^{13}\text{C}$  assignments of **1** and **2** were determined by DEPT, COSY, and HMQC experiments. The conformations of the 17 $\alpha$ -phenylvinyl substituents of **1** and **2** are of interest because of their differing receptor binding affinities and effects in MCF-7 cell proliferation assays. A statistical method of evaluating contributing conformers of **1** and **2** from predicted  $^{13}\text{C}$  shifts of possible structures correlated quite well with conformational conclusions derived from the NOE data. The 17 $\alpha$  substituents of **1** and **2** apparently have a similar conformational preference in solution, suggesting that **1** and **2** could occupy a similar receptor volume.

## Introduction

As part of our efforts to develop more effective therapeutic agents for the treatment of breast cancer, we undertook the designing of (17 $\alpha$ , 20E)-(X-phenyl)vinyl estradiol compounds that can potently and selectively block the interaction of estradiol with its target receptor to impart the desired biological effect. In MCF-7 cell proliferation assay studies, we observed that (17 $\alpha$ , 20E)-(p- $\alpha$ ,  $\alpha$ ,  $\alpha$ -trifluoromethylphenyl)vinyl estradiol (**1**) antagonizes estradiol stimulation and is a very weak agonist with a relative binding affinity (RBA) of 7, compared to the native ligand estradiol at 100.<sup>1</sup> In contrast, under the same assay conditions, (17 $\alpha$ , 20E)-(o- $\alpha$ ,  $\alpha$ ,  $\alpha$ -trifluoromethylphenyl)vinyl estradiol (**2**) was a full agonist with an RBA of 71. We examine here whether *ortho* vs. *para* placement of the substituent in these *E*-isomers could produce a difference between the preferred conformations of **1** and **2** that could account for their distinguishable biological responses and varying binding affinity. In particular, if the conformational preferences differ and that difference were to be preserved upon binding to the receptor, it could affect the conformational mobility of the key helix-12 of the ligand binding domain (LBD) of the estrogen receptor (ER).<sup>2</sup>

Recently we showed that the placement of a substituent in the *ortho* or *para* position in some *Z*-isomers, i.e., (17 $\alpha$ , 20Z)-phenylvinyl estradiols did affect the conformational equilibrium of the 17 $\alpha$  side chain.<sup>3</sup> In that study, (17 $\alpha$ , 20Z)-(p-methoxyphenyl)vinyl estradiol and (17 $\alpha$ , 20Z)-(o- $\alpha$ ,  $\alpha$ ,  $\alpha$ -trifluoromethylphenyl)vinyl estradiol were found to exist in similar conformational equilibria which suggested they would likely occupy a similar receptor volume. These results were consistent with their similar RBA values of 20 and 23. In contrast, (17 $\alpha$ , 20Z)-(o-hydroxymethylphenyl)vinyl

estradiol, which had an RBA of 140, was found to exist in a different conformational equilibrium in terms of the distribution of structures varying in the angles of rotation about the bonds to the vinyl group. These results suggested that, in addition to position and electronic effects of the substituent, the preferred conformations of the  $17\alpha$  substituent of Z-compounds might account for some variation of the RBA values.

In this report, we present a conformational study of the 20E-isomers, **1** and **2**, using NMR and computational methods, to determine whether differences in the preferred conformation of **1** and **2** might also account for their distinguishable biological responses and binding affinities.

The key conformational feature to establish for **1** and **2** is the orientation of the  $17\alpha$  substituent relative to the steroid skeleton. In this study, we use molecular mechanics calculations to generate a set of possible conformations. Two types of NMR data are used in conjunction with the predicted conformations to evaluate which conformations are populated in solution. One approach is to use  $^{13}\text{C}$  chemical shifts in a comparison with shifts predicted for each of the geometries generated from the molecular mechanics calculations. The predicted  $^{13}\text{C}$  shifts come from empirically scaled GIAO (gauge including atomic orbitals) shielding calculations.<sup>4</sup> The other approach is to compare  $^1\text{H}$ - $^1\text{H}$  nuclear Overhauser effects established in one- and two-dimensional experiments, 1D and 2D NOESY, with predicted interatomic distances.

## Experimental

The syntheses and biological data of compounds **1** and **2** have been described elsewhere.<sup>5</sup>  $^1\text{H}$  NMR data were recorded at  $25^\circ\text{C}$  for 5-8 mg samples dissolved in acetone- $d_6$  in 5 mm NMR tubes using a Varian Unity 500 MHz NMR spectrometer

equipped with a 5 mm Varian inverse probe. DEPT and  $^{13}\text{C}$  experiments were obtained on a Varian Mercury instrument at 75 MHz.

$^1\text{H}$  spectra were obtained with a spectral width (SW) of 8 kHz, a  $67^\circ$  pulse flip angle, a 1.7 s acquisition time (AT), a 2 s relaxation delay (RD) and digitized with 32768 points giving a digital resolution (DR) of 0.488 Hz per point. Chemical shifts were referenced to the residual  $^1\text{H}$  signal of acetone- $d_6$ .

$^1\text{H}$ -decoupled  $^{13}\text{C}$  spectra were recorded with a 18856 Hz SW, a  $60^\circ$  pulse flip angle, a 2 s RD and digitized into 65536 points to give a digital resolution of 0.575 Hz per point.

HMQC<sup>6</sup> experiments for single bond  $^1\text{H}$ ,  $^{13}\text{C}$  chemical shift correlation spectra utilized the BIRD sequence to suppress unwanted signals and GARP<sup>7</sup>  $^{13}\text{C}$  decoupling. Two sets of 256 time increments were obtained in the phase-sensitive mode with 32 transients obtained per time increment and a 2 s RD. The final matrix was processed with Gaussian functions.

COSY<sup>45</sup><sup>8</sup> experiments were performed with 8 scans for each of 200 increments in  $F_1$ , 2048 data points in  $F_2$  and a relaxation delay of 2.0 s. The final matrix was symmetrized and processed with sine-bell exponential multiplication.

NOESY<sup>9</sup> experiments were performed with 32 scans for each of 256  $F_1$  increments, 2048 data points in  $F_2$ , with a relaxation delay of 2.0 s and a mixing time of 0.500 s. The final matrix was not symmetrized, but was processed with Gaussian weighing functions.

1D NOESY<sup>10</sup> spectra were obtained using a spectral width of 5000 Hz and 20500 points giving a digital resolution of 0.490 Hz per point, a mixing time of 0.500 s, a RD of



2.0 s, and a AT of 1.7 s. A Gaussian shaped pulse was used for selective irradiation.

## RESULTS AND DISCUSSION

### $^1\text{H}$ and $^{13}\text{C}$ Assignments

The  $^1\text{H}$  NMR spectra of **1** and **2** in acetone- $d_6$  (Figures 1(a) and 2(a)) exhibit very little chemical shift dispersion in the low frequency spectral regions (1.2-2.5 ppm), precluding straightforward  $^1\text{H}$  assignment even at 500 MHz. However,  $^1\text{H}$  signals were assigned via application of HMQC and COSY techniques. The first step was to make  $^{13}\text{C}$  shift assignments, based on our earlier studies of several  $17\alpha$ -substituted estradiols,<sup>3</sup> DEPT experiments, and theoretical shielding calculations (see below). Then, geminal proton resonances were identified and all proton signals were correlated with directly attached carbons via an HMQC experiment. COSY experiments confirmed the initial assignments made by the HMQC experiment but did not, of course, distinguish between  $\alpha$  and  $\beta$  hydrogens in a given methylene group. This distinction was readily achieved by 1D NOESY experiments (Figure 1(b) and 2(b)). Using a Gaussian pulse, selective irradiation of the protons of the methyl group enhances protons on the  $\beta$  face of the C and D rings, viz.,  $11\beta$ ,  $12\beta$ ,  $15\beta$ ,  $16\beta$ , and H8. Table 1 lists the complete assignments of the  $^1\text{H}$  and  $^{13}\text{C}$  signals of **1** and **2**.

### Theoretical Carbon Chemical Shifts and Conformational Determination

The predicted low energy conformers of **1** and **2** (Figures 3 and 4) were generated using the MM3<sup>11</sup> force field through conformational searching by a previously described method.<sup>3</sup> The key dihedral angles are listed in Table 2 for the lowest energy conformers,

**1a-1c** and **2a-2h**, with energies within 1.6 kcal of the lowest energy conformers for **1** and **2**, respectively.

As the MM3 calculations show, significant changes in the  $17\alpha$  side chain conformation result in only minor energy differences. Most of the low energy conformers are within 1 kcal of the lowest energy conformer, making any conformational determination based purely on MM3 energy predictions unreliable. In MMX<sup>12</sup> and MM3 force fields, driving the dihedral angle C21-C20-C17-C13 shows a very shallow energy surface from  $85^\circ$  to  $165^\circ$ . In this region, discrete changes in the orientation of the phenyl to the vinyl group yielded numerous minima using either MM3 or MMX. Conformers **1a** and **1c** were kept as minima since they represent the upper and lower dihedral range of this shallow surface.

Conclusions regarding the preferred  $17\alpha$  side chain conformation of **1** and **2** were approached by applying a statistical method of determining contributing conformers from predicted  $^{13}\text{C}$  chemical shifts,  $\delta_{\text{pred}}$ , of MM3 determined conformers.<sup>3,13</sup> The  $\delta_{\text{pred}}$  were calculated for each MM3-predicted conformer by empirically scaling GIAO-calculated absolute shieldings,  $\sigma$ , obtained at the B3LYP/3-21G level with heteroatoms augmented at the 6-31+G\* level.<sup>4</sup> All shielding calculations were carried out with the Gaussian98 program.<sup>14</sup> Tables 3 and 4 list the  $\delta_{\text{pred}}$  of each MM3 conformer and the assigned experimental  $^{13}\text{C}$  chemical shifts,  $\delta_{\text{exp}}$ .

In this statistical method, the predicted  $^{13}\text{C}$  shifts of the C and D rings of all MM3 conformers of **1** and **2** were in each separate case treated as independent variables in a multiple independent variable regression analysis of the corresponding experimental data.<sup>15</sup>

The predicted  $^{13}\text{C}$  shifts of the A and B rings of all reasonable conformers of **1** and **2** were not used in this statistical analysis since all remain the same within 0.5 ppm regardless of the conformer. In contrast, while the shift differences are still relatively small,<sup>4</sup> most carbons in the C and D rings of **1** and **2** display larger than 1 ppm shift differences depending on the geometry. The regression analysis yielded fractional populations as the fitting parameters. All standard errors and confidence levels of the regression analysis were estimated using the Bootstrapping method.<sup>16</sup>

The results and corresponding estimates of uncertainties (standard errors) are listed in Table 5. Both **1** and **2** were found to have a major conformer, **1b**,  $72 \pm 32\%$ , and **2e**,  $65 \pm 33\%$ . Minor conformers are also indicated for each: **1a**,  $13 \pm 29\%$ , and **1c**,  $15 \pm 28\%$ ; and **2c**,  $33 \pm 18\%$ , and **2h**,  $2 \pm 22\%$ . It is important to note that the large corresponding standard errors make conclusions on the presence of minor conformers unreliable.

The success of this statistical approach for deducing percentage populations of participating conformations in fast exchange depends upon several conditions. Obviously, the molecular mechanics calculations must correctly identify all the possible contributing conformations and represent their conformations appropriately. The chemical shift calculations must also be reasonably accurate; confidence in this regard is engendered by the close match of predicted chemical shifts to experimental shifts at the positions that do not have conformationally dependent shifts. Finally, there must be substantial differences in chemical shifts among the possible conformations and unique combinations of chemical shifts for each. The large uncertainties in the statistical analyses of **1** and **2** shown in Table

5 arise from the relatively small differences in predicted shifts among the different conformers. The largest variations in the C and D rings are predicted to occur at C16 and C14. Thus, in this case it is sensible to take the statistical results only as a qualitative indicator of the major conformers rather than as a meaningful population analysis.

## NOESY Studies

The solution state conformations of the 17 $\alpha$  side chain of **1** and **2** were also investigated by measuring NOE intensities between the vinyl protons and the aliphatic  $^1\text{H}$  of the C and D ring. The 2D NOESY of **1** and **2** reveal a similar pattern of NOE cross peaks and intensities between H20 and H21 with the aliphatic protons 12 $\alpha$ , 12 $\beta$ , H14, 15 $\alpha$ , 16 $\alpha$ , and 16 $\beta$  (Figure 5). Selective 1D NOESY experiments of H20 and H21 allow a more detailed inspection of the NOE intensities (Figures 1(c) and 2(c), (d)). Table 6 and 7 summarize and compare the intensities of the observed NOE signals with expected NOE's based on H-H distances in all predicted low energy conformers of **1** and **2**.

The NOE data for **1** and **2** suggest a similarly preferred orientation of the 17 $\alpha$  vinyl side chain. The absence of an observable NOE between H21 and 12 $\alpha$  rules out conformers, **1c**, **2a**, **2b**, **2c**, **2g**, and **2h**, as contributing conformers, precluding most of the minima observed in the shallow energy surface range of 85° to 165° for dihedral C21-C20-C17-C13. The observable NOE's between H21 and H14, 15 $\alpha$ , and 16 $\alpha$  are consistent with **1b** and **2e**. The presence of the extended conformers, **1a**, **2d**, and **2f** are evident from the weak enhancements of 15 $\alpha$  and 16 $\alpha$  upon irradiation of H20.

## Conclusions

The NOE data of **1** and **2** and the statistical analysis of  $^{13}\text{C}$  chemical shifts are both

consistent with a preferred orientation of the  $17\alpha$  vinyl side chain. The findings from the multiple independent variable linear regression analyses of the  $^{13}\text{C}$  data of **1** and **2**, that conformers **1b** and **2e** are the major conformers, are compatible with the identity of the major conformers favored by NOE data. Conformer **1b** has the vinyl group turned under the D-ring, with the C-21 hydrogen projecting toward C15, and with the C-21 phenyl only slightly twisted out of the plane of the double bond. Conformer **2e** has the double bond disposed in a similar fashion, but has a greater torsion between the double bond and phenyl ring, that allows the ortho substituent to avoid A-strain and to project out from under the D-ring.

The statistical method and the NOE data do not agree on the identity of the minor conformers. For **1**, the regression analysis predicts a 12% population of the extended conformer **1a** that is consistent with the NOE data. However, the minor populations of **1c** and **2c** that appear in the statistical analyses are inconsistent with NOE data that instead favor **2d** and **2f** as minor conformers. The disagreements can be attributed to the limitation of the regression analysis due to the small  $^{13}\text{C}$  shift differences in the C and D ring among most of the MM3 predicted conformers. However, the ability of the regression analysis, based on predicted  $^{13}\text{C}$  shifts, to identify the same major conformer as identified by NOE data demonstrates that this approach to interpretation of chemical shift data is a useful complement to more common methods of conformational analysis.

This study reveals that the  $17\alpha$  substituents of **1** and **2** have a similar preferred orientation in reference to the steroidal skeleton. This similarity in solution conformations of **1** and **2** suggests that there are no inherent restrictions with respect to occupying a

similar receptor volume other than positional placement of the substituent. Other influences such as electronic effects may also play roles in the differing biological responses and RBA values of **1** and **2**. It is interesting to note the apparent absence of a similar range of conformational structures for the 20E-isomers **1** and **2** compared to the 20Z-isomers previously examined,<sup>3</sup> despite the predictions of favorable energies for other conformers in MM3 calculations. However, with only minor energy differences between conformers, this could be a case where the solution conformational preference could be overridden by interactions with the receptor.

### Acknowledgements

We thank Dr. Roger Kautz for valuable assistance concerning the NMR experiments. This work has been supported in part by PHS award R01-CA81<sup>049</sup> and a grant<sup>5</sup> from the U.S. Army DAMD17-99-1-9333, and <sup>DAMD</sup> 17-00-1-0324.<sup>^</sup>

## References


1. Hanson RN, Lee CY, Friel C, DeSombre ER, Hughes A. *Steroids* (Submitted).
2. (a) Brzozowski AM, Pike AC, Dauter Z, Hubbard RE, Bonn T, Engstrom O, Ohman L, Greene GL, Gustafson JA. *Nature* 1997; **389**: 753; (b) Shiau AK, Barstad D, Loria PM, Cheng L, Kushner PJ, Agard DA, Greene GL. *Cell* 1998; **95**: 927.
3. Sebag AB, Friel CJ, Hanson RN, Forsyth DA. *J. Org. Chem.* 2000; **65**: 7902..
4. Forsyth DA, Sebag AB. *J. Am. Chem. Soc.* 1997; **119**: 9483.
5. Lee CY, Hanson RN. *Tetrahedron* 2000; **56**: 1623.
6. Bax A, Subramanian S. *J. Magn. Reson.* 1986; **65**: 565.
7. Shaka AJ, Barker PS, Freeman R. *J. Magn. Reson.* 1985; **64**: 547.
8. Bax A, Freeman R, Morris GA. *J. Magn. Reson.* 1981; **42**: 164.
9. Jeener J, Meier BH, Bachmann P, Ernst RR. *J. Chem. Phys.* 1979; **71**: 4546.
10. Kessler H, Oschkinat H, Griesenger C, Bermel W. *J. Magn. Reson.* 1986; **70**: 106.
11. (a) Allinger NL, Yuh YH, Lii JH. *J. Am. Chem. Soc.* 1989; **111**: 8551; (b) MM3(94) in Alchemy2000, Tripos, Inc.: St. Louis, MO.
12. (a) Gajewski JJ, Gilbert KE, McKelvie H. In *Advances in Molecular Modeling*; Liotta, D (ed.), JAI Press: Greenwich, CT, 1990; Vol. 2. (b) PCMODEL, V.6.0, Serena Software: Box 3076, Bloomington, IN.
13. Sebag AB, Forsyth DA, Plante MA. *J. Org. Chem.* (Submitted).
14. Gaussian 98, Revision A.3, Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Zakrzewski VG, Montgomery, Jr. JA, Stratmann RE, Burant JC,

Dapprich S, Millam JM, Daniels AD, Kudin KN, Strain MC, Farkas O, Tomasi J, Barone V, Cossi M, Cammi R, Mennucci B, Pomelli C, Adamo C, Clifford S, Ochterski J, Petersson GA, Ayala PY, Cui Q, Morokuma K, Malick DK, Rabuck AD, Raghavachari K, Foresman JB, Cioslowski J, Ortiz JV, Stefanov BB, Liu G, Liashenko A, Piskorz P, Komaromi I, Gomperts R, Martin RL, Fox DJ, Keith T, Al-Laham MA, Peng CY, Nanayakkara A, Gonzalez C, Challacombe M, Gill PMW, Johnson B, Chen W., Wong MW, Andres JL, Gonzalez C, Head-Gordon M, Replogle ES, Pople JA, Gaussian, Inc., Pittsburgh PA, 1998.

15. SPSS Base 10.0, SPSS Inc.: Chicago, IL, 1999.

16. Mooney CZ, Duval RD. *Bootstrapping*, Sage Publications, Inc.: Newbury Park, CA, 1993.





## Figure Captions


**Figure 1.** (a) Low frequency spectral region of the 500 MHz  $^1\text{H}$  NMR spectra of **1** in acetone- $\text{d}_6$ . Equivalent spectral regions of the 500 MHz 1D NOESY spectra (500 ms mixing time) of **1** obtained by selective irradiation of the C18 methyl (b), and H20/H21 (c) using a Gaussian pulse. Spectra (b) and (c) are 4x the vertical scale of (a). Overlap of H20 and H21 inhibited selective irradiation of each proton.


**Figure 2.** (a) Low frequency spectral region of the 500 MHz  $^1\text{H}$  NMR spectra of **2** in acetone- $\text{d}_6$ . Equivalent spectral regions of the 500 MHz 1D NOESY spectra (500 ms mixing time) of **2** obtained by selective irradiation of the C18 methyl (b), H20 (c), and H21 (d) using a Gaussian pulse. Spectra (b), (c), and (d) are 4x the vertical scale of (a).


**Figure 3.** MM3-predicted geometries for the most stable conformers of **1**.

**Figure 4.** MM3-predicted geometries for the most stable conformers of **2**.


**Figure 5.** Spectral regions of the 500 MHz 2D NOESY spectrum of (a) **1** and (b) **2** obtained with a mixing time of 500 ms. The NOE connectivities are indicated.














1a






1b




1c







(a)



(b)



**Table 1.**  $^1\text{H}$  and  $^{13}\text{C}$  Chemical Shifts for **1** and **2**

$^1\text{H}$	<b>1</b>	<b>2</b>	$^{13}\text{C}$	<b>1</b>	<b>2</b>
1	7.09	7.09	1	126.4	126.4
2	6.58	6.59	2	112.9	112.9
4	6.52	6.54	3	155.3	155.2
6 $\alpha$	2.75	2.78	4	115.3	115.3
6 $\beta$	2.80	2.81	5	137.7	137.5
7 $\alpha$	1.32	1.34	6	30.0	29.9
7 $\beta$	1.92	1.92	7	27.7	27.7
8	1.43	1.54	8	40.1	40.0
9	2.10	2.10	9	44.0	44.0
11 $\alpha$	2.26	2.28	10	131.3	131.3
11 $\beta$	1.46	1.43	11	26.7	26.6
12 $\alpha$	1.42	1.50	12	32.9	32.8
12 $\beta$	1.68	1.69	13	47.8	47.8
14	1.70	1.71	14	49.5	49.4
15 $\alpha$	1.86	1.84	15	23.5	23.5
15 $\beta$	1.52	1.50	16	37.0	36.9
16 $\alpha$	2.04	2.04	17	83.6	83.7
16 $\beta$	2.06	2.06	18	14.2	14.1
CH <sub>3</sub>	1.02	1.01	20	140.0	141.8
20	6.74	6.65	21	125.3	122.9
21	6.85	7.0	22	142.1	137.8
23	7.69	N/A	23	127.0	127.6
24	7.64	7.70	24	125.6	125.7
25	N/A	7.61	25	128.9	132.6
26	7.64	7.44	26	125.6	127.2
27	7.69	7.82	27	128.2	128.0
N/A	N/A	N/A	CF <sub>3</sub>	125.4	125.7

**Table 2.** Relative Energies and Key Dihedrals of Predicted Conformers of  
1 and 2 Using MM3

Conformers	C13-C17-C20-C21	C20-21-22-23	Relative Energies (kcal/mol)
1a	161	-169	0
1b	-96	18	0.30
1c	89	158	0.32
2a	87	-145	0
2b	95	151	0.06
2c	89	-56	0.92
2d	149	144	1.25
2e	-99	55	1.26
2f	162	-148	1.33
2g	-65	-49	1.53
2h	-95	-68	1.59

**Table 3.** Experimental and Predicted  $^{13}\text{C}$  Chemical Shifts (ppm) of  
Predicted Conformers of **1** Using B3LYP/3-21G(X,6-31+G\*)//MM3

Calculations

Carbon	1a	1b	1c	expt
C1	127.6	127.2	127.3	126.4
C2	113.2	113.1	113.1	112.9
C3	153.3	153.1	153.0	155.3
C4	116.1	115.8	115.7	115.3
C5	136.4	136.0	135.9	137.7
C6	31.2	30.8	30.7	30.0
C7	28.2	28.3	28.3	27.7
C8	40.3	39.8	39.5	40.1
C9	44.2	43.9	43.9	44.0
C10	131.6	131.5	131.8	131.3
C11	28.3	28.1	28.2	26.7
C12	31.4	31.8	32.4	32.9
C13	46.5	48.1	48.1	47.8
C14	50.2	48.0	47.9	49.5
C15	25.7	26.2	26.4	23.5
C16	45.9	36.8	39.2	37.0
C17	83.0	84.9	84.1	83.6
C18	15.2	16.6	15.5	14.2
C20	149.2	144.9	143.4	140.0
C21	133.9	131.6	131.3	125.3
C22	137.9	138.6	137.6	142.1
C23	121.6	122.4	122.0	127.0
C24	127.5	127.6	127.5	125.6
C25	132.2	131.7	131.8	128.9
C26	128.0	127.8	127.8	125.6
C27	129.3	128.5	129.2	128.2
C28	130.8	130.9	130.8	125.4

**Table 4.** Experimental and Predicted  $^{13}\text{C}$  Chemical Shifts (ppm) ofPredicted Conformers of **2** Using B3LYP/3-21G(X,6-31+G\*)//MM3

Calculations

Carbon	2a	2b	2c	2d	2e	2f	2g	2h	expt
C1	127.4	127.6	127.3	127.5	127.5	127.4	127.6	127.6	126.4
C2	113.1	113.1	113.0	113.1	113.1	113.1	113.1	113.0	112.9
C3	153.1	152.9	153.0	153.0	153.0	153.1	153.1	152.9	155.2
C4	115.8	115.5	115.8	115.8	115.7	115.9	115.8	115.7	115.3
C5	136.2	135.8	136.1	136.1	136.1	136.3	136.2	136.2	137.5
C6	31.1	30.8	30.7	31.1	31.1	30.9	31.1	31.1	29.9
C7	28.3	28.4	28.1	28.3	28.3	28.2	28.4	28.3	27.7
C8	39.9	39.7	40.1	39.9	39.5	40.1	40.3	40.0	40.0
C9	44.3	44.4	44.2	44.3	44.3	44.3	44.5	44.2	44.0
C10	131.6	132.1	131.6	131.6	132.0	131.5	131.7	132.1	131.3
C11	28.3	28.5	28.5	28.2	28.3	28.2	28.4	28.4	26.6
C12	33.0	33.8	33.1	31.9	32.2	31.9	29.4	30.5	32.8
C13	48.1	48.6	48.0	47.7	47.2	47.2	48.6	47.6	47.8
C14	51.5	51.1	50.9	49.6	48.3	49.7	50.8	47.6	49.4
C15	26.0	26.5	26.0	25.8	26.4	25.6	26.8	26.1	23.5
C16	39.6	42.4	39.2	39.6	38.9	41.6	39.7	37.6	36.9
C17	83.6	83.9	84.6	84.0	83.6	84.0	81.9	85.5	83.7
C18	16.3	15.0	16.1	16.3	15.2	16.3	18.3	16.9	14.1
C20	146.2	146.7	151.2	146.3	151.1	147.5	149.6	151.2	141.8
C21	130.7	133.3	130.1	129.9	225.7	132.5	128.7	131.7	122.9
C22	137.2	137.5	140.8	138.6	140.4	138.7	140.7	140.6	137.8
C23	126.8	126.4	128.1	127.3	128.4	126.6	128.7	129.0	127.6
C24	131.9	131.9	130.7	132.1	130.9	131.9	130.9	131.0	125.7
C25	130.6	130.6	130.9	130.9	131.0	130.7	130.9	130.8	132.6
C26	127.8	127.6	129.5	127.7	129.5	127.7	129.3	129.0	127.2
C27	130.5	130.1	131.4	130.1	131.2	130.0	131.0	130.5	128.0
C28	126.8	126.5	126.0	126.3	126.2	126.6	126.3	126.1	125.7

**Table 5.** Summary of the Multiple Independent Variable Regression

Analysis<sup>a</sup> of the Calculated <sup>13</sup>C Shifts of Predicted Conformers of 1 and 2

Conformer	Estimate (%)	Standard Error (%)
1a	13	29
1b	72	32
1c	15	28
2a	0	14
2b	0	13
2c	33	18
2d	0	18
2e	65	33
2f	0	30
2g	0	4
2h	2	22

<sup>a</sup> Constraints: Each conformer is greater than or equal to 0 %. Each of the conformer sets **1a-1c** and **2a-2h** total to 100 %.


**Table 6.** Summary and Comparison of Observed NOE Enhancements with Expected NOE Intensities<sup>a</sup> for Predicted Conformers of **1**

<b>Irradiated</b>	<b>Enhanced</b>	<b>1a</b>	<b>1b</b>	<b>1c</b>	<b>Expt</b>
H20	12 $\alpha$	s	s	w	<b>s</b>
H20	H14	s	s	s	<b>s</b>
H20	15 $\alpha$	w	n	w	<b>w</b>
H20	16 $\alpha,\beta$	w	n	w	<b>w</b>
H21	12 $\alpha$	n	n	s	<b>n</b>
H21	H14,12 $\beta$	n	s	w	<b>w</b>
H21	15 $\alpha$	n	w	n	<b>w</b>
H21	16 $\alpha,\beta$	n	s	n	<b>s</b>


a. Expectations of strong (s), weak (w), and no (n) NOE enhancements correspond to H-H distances of 0 - 2.99; 3.0 - 4.99; and > 5 Å.

---

*The Philip S. Portoghese  
Symposium in  
Medicinal Chemistry*



*August 23-24, 2001  
2-530 Moos Tower  
University of Minnesota*



*Philip S. Portoghese, Ph.D.*



---

Philip S. Portoghese, Ph.D., is a University Distinguished Professor of Medicinal Chemistry at the University of Minnesota. A graduate of the University of Wisconsin Medicinal Chemistry department, he has served on the faculty at Minnesota since 1961. Dr. Portoghese is a member of the editorial or editorial advisory boards of many journals and scientific organizations, and has been the Editor-in-Chief of the *Journal of Medicinal Chemistry* since 1972. He has received honorary doctorate degrees from the University of Catania, Italy and the Royal Danish School of Pharmacy. He is the recipient of many awards including the A.Ph.A. Research Achievement Award (1980), the Volwiler Award (1984), the ACS Medicinal Chemistry Award (1990), the AAPS Research Achievement Award in Medicinal Chemistry (1990), the Nathan Eddy Award (1991), the ACS Edward Smissman Award (1991), the Rho Chi Pharmacy Honor Society Award (1999), the Italian Chemical Society Oak & Tulip Award (1999), and the ACS Alfred Burger Award (2000). He is a fellow of the Academy of Pharmaceutical Sciences (1974), AAAS (1986), AAPS (1986) and College on Problems of Drug Dependence (1990).

Prof. Portoghese has presented numerous lectures at national and international conferences, universities and industry, has authored over 315 publications and has trained approximately 95 students and postdoctoral research associates. His research interests are in the area of drug design with specific contributions to the area of opioid chemistry. The impact of his work on the field of drug design and receptor-ligand interactions is exemplified by him being listed as one of the most cited authors in the area of xenobiotics for the period 1981-1992.

On the occasion of his 70<sup>th</sup> birthday, we would like to celebrate Phil's outstanding contributions not only as a preeminent scholar in medicinal chemistry, but also in his teaching and mentoring of scientists, as well as his outstanding service as Editor of the *Journal of Medicinal Chemistry*. He has uniquely contributed to the quality and reputation of the Department of Medicinal Chemistry, the College of Pharmacy, and the University of Minnesota.

# PROGRAM

August 23, 2001  
2-530 Moos Tower

8:30 Registration

9:00 Welcome and Introduction

Yusuf Abul-Hajj, Ph.D., Professor and Head  
*Department of Medicinal Chemistry, University of Minnesota*

Marilyn Speedie, Ph.D., Dean  
*College of Pharmacy, University of Minnesota*

MODERATOR: Yusuf Abul-Hajj, Ph.D., *University of Minnesota*

9:15 "SPECIFIC GABA<sub>A</sub> RECEPTOR LIGANDS: FROM MUSHROOM  
CONSTITUENT THROUGH CLINICALLY ACTIVE DRUG TO  
RATIONAL DRUG DESIGN"

Povl Krosgaard-Larsen, Ph.D., *Royal Danish School of Pharmacy*

9:55 "DOPAMINE RECEPTORS IN THE CENTRAL NERVOUS  
SYSTEM: TARGETS FOR MEDICINAL CHEMISTS PAST,  
PRESENT AND FUTURE"

John L. Neumeyer, Ph.D., *Harvard University*

10:35-11:00 Break

11:00 KEYNOTE ADDRESS

Ralph Hirschmann, Ph.D., *University of Pennsylvania*

12:00-1:30 Lunch Break

MODERATOR: Rick Wagner, Ph.D., *University of Minnesota*

1:30    ***"DESIGN AND SYNTHESIS OF PEPTIDASE  
INHIBITORS. WHAT CHALLENGES REMAIN?"***

Daniel H. Rich, Ph.D., *University of Wisconsin*

2:10    ***"DESIGN, SYNTHESIS AND EVALUATION OF NOVEL  
STEROIDAL ANTIESTROGENS FOR THE TREATMENT OF  
HORMONE RESPONSIVE BREAST CANCER"***

Robert N. Hanson, Ph.D., *Northeastern University*

2:50-3:20      **Break**

3:20    ***"G-QUADRUPLEXES AND ASSOCIATED GENE TARGETS  
FOR DRUG DESIGN"***

Laurence H. Hurley, Ph.D., *University of Arizona*

4:00    ***"BENZODIAZEPINE CCK-A RECEPTOR AGONISTS"***

Elizabeth E. Sugg, Ph.D., *GlaxoSmithKline Company*

4:40    ***"2(3H)-BENZOTHAZOLONES, AN INEXHAUSTIBLE  
SOURCE OF INSPIRATION FOR AN ACADEMIC  
MEDICINAL CHEMIST. APPLICATION TO THE DESIGN  
OF MIXED AFFINITY LIGANDS FOR 5-HT SUB-CLASSES"***

Jacques Poupaert, Ph.D., *University of Louvan, Belgium*

6:30    **Reception – Cash Bar**  
McNamara Alumni Center, A.I. Johnson Great Room

7:30    **Dinner – Pre-registered guests only**  
McNamara Alumni Center, A.I. Johnson Great Room

August 24, 2001  
2-530 Moos Tower

MODERATOR: David Ferguson, Ph.D., *University of Minnesota*

8:30 "OPIOID RECEPTORS: STRUCTURAL INSIGHTS INTO  
RECEPTOR FUNCTION AND LIGAND INTERACTION FROM  
MOLECULAR MODELING"

M. Germana Paterlini, Ph.D., *University of Minnesota*

9:00 "EVOLUTION OF THE 3,4-DIMETHYL-4-PHENYLPYPERIDINE  
OPIOID ANTAGONISTS: FROM DISCOVERY AND  
PHARMACOLOGICAL CHARACTERIZATION TO  
THERAPEUTIC DEVELOPMENT"

Dennis M. Zimmerman, Ph.D., *Lilly Research Laboratory*

9:40 "DEVELOPMENT OF SELECTIVE OPIOID ANTAGONISTS"

F. Ivy Carroll, Ph.D., *Research Triangle Institute*

10:20-10:50 Break

10:50 "EXPLORING THE CHEMISTRY OF PAIN AND ADDICTION  
WITH A NEW PARADIGM."

Victor J. Hruby, Ph.D., *University of Arizona*

11:30 "OPIOID PEPTIDE ANALOGUES AND PEPTIDOMIMETICS AS  
PHARMACOLOGICAL TOOLS AND POTENTIAL DRUGS"

Peter W. Schiller, Ph.D., *Research Institute of Montreal*

12:10-1:30 Lunch Break

MODERATOR: Herbert Nagasawa, Ph.D., *University of Minnesota*

1:30 **"OPIOID RESEARCH TOOLS AND DRUGS FOR THE  
NEW MILLENNIUM"**

Kenner C. Rice, Ph.D., *National Institutes of Health*

2:10 **"DESIGN AND SYNTHESIS OF NOVEL PEPTIDE-BASED  
AFFINITY LABELS FOR OPIOID RECEPTORS"**

Jane V. Aldrich, Ph.D., *University of Maryland*

2:50-3:20      Break

3:20 **"DEVELOPMENT OF POTENT AND HIGHLY SELECTIVE  
SMALL MOLECULE INHIBITORS OF VLA4"**

Francine S. Grant, Ph.D., *Elan Pharmaceuticals*

4:00 **"SELECTIVE MECHANISM-BASED INHIBITION OF THE  
QUINONE-DEPENDENT AMINE OXIDASES"**

Lawrence M. Sayre, Ph.D., *Case Western Reserve University*

4:40 **Symposium Summation**

Yusuf Abul-Hajj, Ph.D.

4:45 **"WORDS OF WISDOM"**

Philip S. Portoghese, Ph.D.

## Graduate Students and Research Associates

Anthony Abatjolgou	Robert Hanson	Sathit Niratisai
Mohamed Abdel-Monem	James Henkel	Shigenori Ohkawa
Jane Aldrich	Jacob Herzig	Sandra Olmsted
Bipin Alreja	Sandor Hosztafi	Germana Paterlini
Jianguo An	Mei Hua	Veronika Phillips
Dimitrios Barbas	Masako Ikeda	Jacques Poupaert
Kevin Bell	Jonathan James	Michael Powers
Eric Berg	Jack Jiang	K. Ramakrishan
Rashmi Bushan	Robert Jones	Michael Rein
MariaLaura Bolognesi	Sheung-Tsam Kam	Thomas Riley
Vijaya Boyapati	Renaud Kiesgen de Richter	James Rodgers
An-Chih Chang	Stacy Kramer	Giuseppe Ronsisvalle
Raymond Conrow	Tushar Kshirsagar	Susan Roseff
David Daniels	Amol Kulkarni	Samir Salib
Victoria Darrow	Didier Lambert	Victoria Sandberg
Barbara Di Giacomo	Dennis Larson	Lourdes Santana
Richard Elliott	Danny Lattin	Lawrence Sayre
Mordechai Erez	Bertrand Le Bourdonnec	Joseph Schoenecker
Mohamed Essawi	Chia-En Lin	Dennis Sepp
Christopher Etienne	Andrew Lipkowski	Shiv Sharma
Mei Fang	Mary Lunzer	William Stevens
Sunan Fang	S. Madusoodanan	Idalia Stark
Xinqin Fang	Keith Maloney Huss	Elizabeth Sugg
Francine Farouz	Sante Martelli	Mumtaz Sultana
Thomas Fitzgerald	Christopher McCurdy	Barbara Taylor
Reynold Francis	Arem Melikian	Vasant Telang
David Fries	Thomas Metzger	Joseph Turcotte
Peng Gao	Adel Mikhail	Robert Whyte
Aaron Garzon	Scott Moe	David Williams
Zeinab Gomaa	Mosad Mohamed	Chul-Bu Yim
William Groutas	Joan Naeseth	Yan Zhang
	Hiroshi Nagase	

We wish to thank these individuals for their contributions:

Joy Loban, Department of Medicinal Chemistry  
Joyce F. Reha, Department of Medicinal Chemistry  
Mary Lunzer, Department of Medicinal Chemistry  
Amy Olson, College of Pharmacy